

THE Ca, K and Mg RATIO IN SOIL AND VINES UNDER EFFECTS OF FERTILIZATION WITH DIFFERENT POTASSIUM FERTILIZER DOSES

Raportul Ca, K și Mg în sol și viță de vie sub efecte de fertilizare cu doze diferite de potasiu îngrășământ

Ličina, V. (1), Marković, N. (1), Trajković, I. (1), Atanacković, Z. (1)

(1) Belgrade University, Faculty of Agriculture, Nemanjina 6, 11080 Belgrade, Serbia, e-mail: zoranata4@yahoo.com

The investigation was carried out in the vineyards of "Radmilovac" on the Faculty of Agriculture experimental station, with cv. Sauvignon blanc grafted to the rootstock Berlandieri x Riparia Kober 5BB. The aim of this research was to determine the effects of potassium fertilizer application on the Ca:Mg and K:Mg ratio in soil and vines organs. In this experiment the following treatments were used: control (without fertilization) and treatments with 50, 100 and 150 kg K₂O/ha 50% KCl potassium fertilizer. The soil type was eutric cambisol. Treated soil was subjected to the detail agrochemical soil analysis, while the soil samples was collected from the depth of 0-30, 30-60, 60-90 and 90-120 cm. Leaves for analysis were collected in august and shoots after pruning. Potassium and magnesium analysis has been by AAS. The level of available potassium was (11,95-14,15 mg/100g of soil), magnesium (20,2-23,7 mg/100 g of soil) and calcium (354-464 mg/100 g of soil). During the first year, Ca:Mg ratio was 5,8-14,4:1, in the second year 5,4-18,5:1 and at third year 4,9-25,2:1. The K:Mg ratio ranged from 0,08-0,21:1 in the first, 0,13-0,29:1 in the second and 0,11-0,21:1 in the third year of study. The Ca:Mg and K:Mg ratio were mostly influenced by 100 and 150 kg K₂O/ha potassium doses. The K:Mg ratio in the leaves and shoots did not change under influence of different potassium fertilizer doses, so that the antagonism between these two elements was not manifested.

Key words: potassium fertilizer, Ca, Mg, K

Cercetare a fost realizată în podgoriile de "Radmilovac", la Facultatea de Agricultură stație experimentală, cu cv. Sauvignon blanc altoit la portaltol Berlandieri x Riparia Kober 5BB. Scopul acestui studiu a fost de a determina efectele aplicării îngrășămintelor pe potasiu Ca:Mg și K:Mg raportul în sol și organe viță de vie. În acest experiment tratamente au fost utilizate următoarele: de control (fără fertilizare) și tratamente cu 50, 100 și 150 kg K₂O/ha de potasiu îngrășămintă 50% KCl. Tipul de sola fost cambisol eutric. Sol tratați a fost supus analizei solului detaliu agrochimice, în timp ce probele de sol au fost colectate de la adâncimea de 0-30, 30-60, 60-90 și 90-120 cm. Frunze de analiză au fost colectate în august și lăstarii după tăiere. Analiza potasiu și magneziu a fost facute cu AAS. Nivelul de potasiu a fost disponibil (11,95-14,15 mg/100 g de sol), magneziu (20,2-23,7 mg/100 g de sol) și de calciu (354-464 mg/100 g de sol). Pe parcursul primului an, Ca:Mg raportul a fost 5,8-14,4:1, în al doilea an 5,4-18,5:1 și la treilea an 4,9-25,2:1. K:Mg raportul au variat de la 0,08-0,21:1 în primul an 0,13-0,29:1, în al doilea 0,11-0,21:1 și în al treilea an de studiu. Ca:Mg și K:Mg raportului au fost în mare parte influențate de 100 și 150 kg K₂O/ha doze de potasiu. K:Mg raportului în frunzele și lăstarii nu a schimbat sub influența unor doze diferite de potasiu îngrășămintă, astfel că antagonismul dintre aceste două elemente nu sa manifestat.

Cuvinte cheie: îngrășămintă potasiu, Ca, Mg, K

INTRODUCTION

Balanced nutrition and fertilization are essential components of the growing technology in vineyards in terms of achieving the required yields and quality of grapes. Potassium fertilizers are soluble in water, act now and quickly adsorbed by plants. In the vineyard practice, recommendation is to use liquid potassium fertilizers in the planting and in the regular production of grapes, especially if it is known the yield of 1000 kg present 10 kg of pure potassium (Nakalamić et al., 2009). Potassium is one of the most mobile elements in plant organs. Unlike nitrogen and phosphorus that are accumulate in reproductive organs, potassium is acumulate in vegetative, such as roots, shoots and leaves (0,2 to 2%). Intensive potassium fertilization and it is increasing adsorption may condition the distortion K:Mg ratio in plant organs. Until this phenomenon usually occurs on soils rich with Mg, where due the increased of K content in soil increases K^+ competition with a lot less frequent Mg^{+2} in the soil solution (Ličina, 2009). The presence of Ca in the soil is important to establish a relationship between certain cations in the soil, which is expressed through mutual ratio between Ca:Mg, Mg:K and Ca:K. This quotient is used as an indicator of plant nutritional level of these elements, especially if it is known that Ca and Mg treat antagonistically in the soil (McLean et al., 1983). Ca:Mg ratio usually varies from 2:1 to 10:1 while the ideal ratio is 4:1 (Conradie, 1994). Garcia et al. (1999) found that an increased content of Ca in the soil causes an increase content of K in petioles and leaf blades, and total acid content in the wine on grapes grown on acidic soils. Ličina et al. (1997) indicate that the high content of potassium in the soil does not be a parameter for the high potassium content in the grapes leaves. This is considered a result of potassium redistribution inside the leaves and other organs (root and stem), where potassium take the other metabolic role. Despite the expressed Mg mobility, because of its deficit or impaired of K:Mg ratio comes to the appearance of necrosis of grape organs and berries (Hlušek et al., 2002). Gluhic et al. (2009) and Zatloukalova et al. (2011) by research are found that the higher Mg content in the grapes organs carry out to distortion of the ideal K:Mg ratio as well as an increase content of K in the organs and a decrease in of Ca content.

MATERIALS AND METHODS

The experiment were carried out in the vineyard of Demonstration Field "Radmilovac" at Faculty of Agriculture in Belgrade on cv. Sauvignon blanc, which was grafted to the rootstock *Berlandieri x Riparia* Kober 5BB. The vineyard is located on the slope of southern exposure. The lines extending in the direction east-southeast. The line spacing is 3 m and the distance between vines in the row is 1m. Training system is double asymmetric cordon (Nakalamić, 1991). The experiment was set up according to the block system with 18 vines arranged in three replications with 6 vines. Immediately before of fertilizers application it is done a detailed agrochemical soil analysis. For the analysis soil is sampled to a depth of 0-30, 30-60, 60-90 and 90-120 cm with the opening of the two profiles between the lines within each treatment. In the experiment was used 50% of KCl in the following doses: 50, 100, 150 kg K_2O /ha and control (without fertilization). With potassium also is added nitrogen (30 kg/ha) and phosphorus (50 kg P_2O_5 /ha). Nitrogen was added in the form of KAN, and phosphorus in the form of triple superphosphate. Leaves for analysis were sampled in August with all the vines in the treatment from the lower, middle and upper shoots. Shoots were sampled after vine cutting within treatment experiment. The data in this paper represent averages of repetitions for each treatment trial. Exchangeable potassium in the soil is determined by the Al-method according to Egner-Riehm, and in leaves and shoots of the wet method of destruction with nitric acid after which the extract was determined by flame photometry (Džamić et al., 1996). Ca and Mg determination in soil and plant material was carried out using AAS.

RESEARCH RESULTS

Before applying the appropriate fertilizer dose by treatments was done detailed agrochemical analysis of soil. For the soil analysis samples were collected from four depths (0-30, 30-60, 60-90 and 90-120 cm), as shown in table 1.

Table 1

Agrochemical properties of the test plots before fertilization with K fertilizers

Depth (cm)	pH		Humus	N	NH ₄	NO ₃	P ₂ O ₅	K ₂ O	Ca	Mg
	H ₂ O	nKCL	%	%	mg/kg	mg/kg	mg/100 g	mg/100 g	mg/100 g	mg/100 g
0-30	7,4	6,5	3,30	0,21	10,5	3,50	15,0	11,95	464	23,2
30-60	7,4	6,5	2,11	0,19	10,5	5,25	8,7	14,15	404	20,2
60-90	6,7	5,4	1,88	0,16	10,0	5,25	7,8	12,75	354	22,1
90-120	6,9	5,7	1,64	0,16	12,1	7,00	8,7	12,25	399	23,7

The results indicate that soil was neutral pH reaction, whose acidity slightly increases with depth. The surface layer of soil pH is 7,4 in H₂O, and the mild acidity (pH 6,7-6,9 in H₂O), represented in layers to a depth of 60-90 cm and 90-120 cm. Percentage of total N ranged from 0,16% to 0,21% and slightly decreases with depth, which this land is classified as high amounts of nitrogen (0,1 to 0,2%), which fits most eutric cambisol in Serbia. Total nitrogen, however, does not correspond with the level of available forms of nitrogen (NH₄+NO₃) which level in the profile is quite high (210 kg N/ha at a depth of 0-120 cm), and was used small doses of nitrogen fertilizer during the research period (30 kg N/ha). The humus content is satisfactory for growing grapes. According to the available phosphorus content, soil in the deeper layers is poor, while the surface layer of high amounts of phosphorus. Potassium contents ranged from 11,95-14,15 mg K₂O/100 g soil, so that this land belongs to the medium amounts of soil potassium. Calcium was the highest in the layer of 0-30 cm (464 mg/100 g soil) and lowest in the layer from 90-120 cm (399 mg/100 g soil). Mg content has not significantly changed in the soil profile.

Table 2

Exchangeable K, Ca and Mg and their relationship to soil samples in the first year of research

Treatment	Profile depth (cm)	K mg/100 g	K mekv/100 g	Ca mg/100 g	Ca mekv/100 g	Mg mg/100 g	Mg mekv/100g	Ca:Mg	K:Mg
Control	0-30	11,95	0,30	388	19,40	23,7	1,95	9,94:1	0,15:1
	30-60	14,15	0,36	478	23,90	21,2	1,74	13,7:1	0,21:1
	60-90	12,75	0,32	399	19,90	27,5	2,26	8,8:1	0,14:1
	90-120	12,25	0,31	354	17,70	36,2	2,98	5,9:1	0,10:1
50 kg/ha	0-30	12,75	0,32	605,5	30,30	35,0	2,88	10,5:1	0,11:1
	30-60	13,60	0,35	461,5	23,10	23,8	1,95	11,8:1	0,18:1
	60-90	14,70	0,37	500,0	25,00	36,2	3,00	8,3:1	0,12:1
	90-120	13,60	0,35	449,0	22,40	36,2	2,98	7,5:1	0,12:1
100 kg/ha	0-30	13,85	0,36	464,0	23,20	32,5	2,67	8,7:1	0,13:1
	30-60	13,60	0,35	404,0	20,20	29,0	2,36	8,6:1	0,15:1
	60-90	14,15	0,36	441,5	22,10	34,0	2,90	7,6:1	0,12:1
	90-120	13,35	0,341	474,0	23,70	50,0	4,11	5,8:1	0,08:1
150 kg/ha	0-30	15,00	0,38	563,0	28,15	23,7	1,95	14,4:1	0,19:1
	30-60	16,10	0,41	536,0	25,33	23,7	1,96	12,9:1	0,20:1
	60-90	14,70	0,37	508,0	25,40	26,2	2,16	11,8:1	0,17:1
	90-120	12,20	0,31	485,0	22,75	30,0	2,47	9,2:1	0,12:1
LSD _{0,05}						0,0450			
LSD _{0,01}						0,0745			

Magnesium as a cation ion exchanger in which the reactions are often replaced by calcium, in relation to potassium is present in greater quantity in the soil and it is reflected

on the K:Mg ratio which is ranged from 0,08-0,21:1 in first year, 0,13-0,29:1 in the second and 0,18:1 in the third research year. This shows that potassium fertilization affected the relationship between these two elements in the soil which is reflected in some variants assessed as statistically significant.

In the first investigation year (table 2) in the profile depth of 30-60 cm, a fertilization treatment with 150 kg K₂O/ha and control showed significant variation in the K:Mg ratio in relation to treatments with 50 and 100 kg K₂O/ha. In profile depth of 90-120 cm K:Mg ratio was more expressed in the fertilization treatments with 150 and 50 kg K₂O/ha compared with the control and the treatment with 100 kg K₂O/ha. Ca:Mg ratio had the highest values in all profile layers in the treatment with 150 kg K₂O/ha (9,2-14,4:1). Other treatments showed less impact on the Ca:Mg ratio changing in the soil where the treatment with 100 kg K₂O/ha had the lowest ratio values.

Table 3

Exchangeable K, Ca and Mg and their relationship to soil samples in the second year of research

Treatment	Profile depth (cm)	K mg/100 g	K mekv/100 g	Ca mg/100 g	Ca mekv/100 g	Mg mg/100 g	Mg mekv/100g	Ca:Mg	K:Mg
Control	0-30	20,0	0,51	548,0	27,1	27,5	2,26	12,0:1	0,22:1
	30-60	19,7	0,50	484,0	24,2	22,5	1,85	13,1:1	0,27:1
	60-90	17,7	0,45	506,5	26,3	27,2	2,26	11,6:1	0,20:1
	90-120	17,8	0,45	325,0	16,2	36,2	2,98	5,4:1	0,15:1
50 kg/ha	0-30	22,5	0,57	669,0	33,4	30,0	2,47	13,5:1	0,23:1
	30-60	21,4	0,54	457,5	22,9	28,7	2,36	9,7:1	0,23:1
	60-90	20,6	0,53	417,5	20,87	36,3	2,96	7,0:1	0,18:1
	90-120	20,8	0,53	415,5	20,8	46,2	3,80	5,5:1	0,14:1
100 kg/ha	0-30	21,1	0,54	736,5	36,8	28,8	2,40	15,3:1	0,22:1
	30-60	20,6	0,52	474,0	23,7	32,5	2,67	8,9:1	0,19:1
	60-90	23,8	0,60	489,0	24,4	36,2	2,98	8,2:1	0,20:1
	90-120	21,6	0,55	511,5	25,6	50,0	4,11	6,2:1	0,13:1
150 kg/ha	0-30	23,3	0,59	764,0	38,2	25,0	2,06	18,5:1	0,29:1
	30-60	21,7	0,55	582,5	29,1	27,5	2,26	12,9:1	0,24:1
	60-90	20,5	0,52	488,0	24,4	28,7	2,36	10,3:1	0,22:1
	90-120	19,4	0,49	471,5	23,6	38,7	3,18	7,4:1	0,15:1
LSD _{0,05}						0,0460			
LSD _{0,01}						0,0763			

In the second year of studies in the profile layer of 0-30 cm fertilization treatment with 150 kg K₂O/ha has significantly higher K:Mg ratio compared to the control and treatment with 100 kg K₂O/ha. Observe with the profile depth values of K:Mg ratio slightly decreasing, and have lower values than the surface layer. That, the profile depth from the 60-90 and 90-120 cm have similar value of K:Mg ratio (0,13-0,22:1) while in the layer of 0-30 cm and 30-60 have higher values approach the values 0,27-0,29:1. Observed all treatments by the profile depth reflected the value of Ca:Mg ratio with increasing depth decreased. The highest values of Ca:Mg ratio (18,5:1) were recorded during treatment with 150 kg K₂O/ha in a profile layer of 0-30 cm (table 3).

Table 4

Exchangeable K, Ca and Mg and their relationship to soil samples in the third year of research

Treatment	Profile depth (cm)	K mg/100 g	K mekv/100 g	Ca mg/100 g	Ca mekv/100 g	Mg mg/100 g	Mg mekv/100g	Ca:Mg	K:Mg
Control	0-30	15,0	0,38	499,0	24,9	27,5	2,25	11,1:1	0,17:1
	30-60	14,7	0,38	441,5	22,1	25,0	2,05	10,8:1	0,18:1
	60-90	12,8	0,33	342,5	17,2	32,5	2,70	6,4:1	0,12:1
	90-120	15,8	0,40	344,0	17,2	39,0	3,20	5,4:1	0,12:1
50 kg/ha	0-30	15,0	0,38	423,0	21,2	24,0	2,00	10,6:1	0,19:1
	30-60	14,7	0,37	440,0	22,	24,0	1,95	11,3:1	0,19:1
	60-90	15,8	0,40	380,0	19,0	37,5	3,10	6,13:1	0,13:1
	90-120	17,5	0,44	357,5	17,9	36,5	3,00	6,00:1	0,15:1
100 kg/ha	0-30	17,2	0,44	400,0	20,0	26,5	2,20	9,1:1	0,20:1
	30-60	17,2	0,44	402,5	20,1	25,5	2,10	9,6:1	0,21:1
	60-90	14,7	0,37	360,0	18,0	39,0	3,20	5,6:1	0,12:1
	90-120	15,3	0,39	331,5	16,6	41,5	3,40	4,9:1	0,11:1
150 kg/ha	0-30	16,7	0,43	616,5	30,8	31,5	2,60	11,8:1	0,16:1
	30-60	16,6	0,42	472,5	23,7	25,0	2,05	11,6:1	0,20:1
	60-90	11,1	0,28	1311,5	65,6	31,5	2,60	25,2:1	0,11:1
	90-120	13,0	0,33	534,0	26,7	35,0	2,90	9,2:1	0,11:1
LSD _{0,05}						3,5018			
LSD _{0,01}						5,807			

In the third year of study (table 4) statistical significance was achieved only in the difference in Ca:Mg ratio in the profile layer of 90-120 cm. Among increase in the value of Ca:Mg ratio in the soil was mostly influenced by treatment with 150 kg K₂O/ha fertilizer where values were the highest compared to other treatments. K:Mg ratio in the third year of investigation was not significantly changed by treatments above than the treatment with 100 kg K₂O/ha where is recorded slightly higher ratio value (0,21:1) in the profile layer of the 0-30 and 30-60 cm.

Table 5

Relations K:Mg in the leaves and shoots during the test period

Treatment	I year		II year		III year	
	Leaf	Shoot	Leaf	Shoot	Leaf	Shoot
Control	6,43:1	11,1:1	5,26:1	6,22:1	4,20:1	4,80:1
50 kg/ha	5,73:1	9,45:1	6,72:1	6,20:1	4,70:1	4,90:1
100 kg/ha	6,26:1	10,0:1	6,65:1	6,20:1	4,17:1	6,44:1
150 kg/ha	6,18:1	8,66:1	6,78:1	8,12:1	4,56:1	6,88:1

The K:Mg ratio in leaves should vary between 3 and 7. Smaller or larger values indicate a deficiency of one nutritional elements (Levy et al., 1964). Among experimental cultivar there was no potassium deficiency in the control and in the treatments with fertilizers, ie. ratio was higher than 3. Also not obvious magnesium deficiency because this quotient was higher than 7. The ratio is not varied under fertilization influence so that with these doses of potassium fertilizers (50-150 kg K₂O/ha) potassium antagonism is not manifested. The analysis of shoots showed that the values of K:Mg ratio were the lowest in the first experiment year in treatment with 150 kg K₂O/ha (8,66:1). In the second and third experimental year there was increasing effect of fertilization with 150 kg K₂O/ha and increased K:Mg ratio in shoots (8,12:1 and 6,88:1) compared to treatment with 50, 100 and 150 kg K₂O/ha and control where the value of K:Mg ratio ranged from 6,20-6,22:1 (in the

second year of investigation) and 4,80-6,66:1 (in the third year of study). The results are shown in table 5.

CONCLUSIONS

Based upon the research can be performed following conclusions:

- Agrochemical analysis showed that soil conditions are favorable for growing grapes and development. With humus content of 3.3% of this soil is considered to be well assured, but the humus content increasing with depth. The level of total nitrogen ranged from 0.16 to 0.21% and is match high. In content of phosphorus in the soil surface layer is characterized as medium provided, while the deeper soil layers is poor in phosphorus;
- The potassium level in soil varied from 11.95 to 14.15 mg K₂O/100 g of soil and that categorize this soil in middle category provided with potassium;
- Soil provide with Ca and Mg were within the optimum;
- Available amounts of Ca (400-600 mg/100 g) and Mg (20-35 g/100 g) provide a normal grape nutrition so Ca:Mg ratio in the soil during the experiment was favorable.
- K:Mg ratio takes, in the first year of experiment values as from 0,08-0,21:1, the second year 0,13-0,29:1 and third year 0,18:1. That indicating that the fertilization with different doses of potassium fertilizer affected the ratio between these two elements in the soil, as in the variants with 100 and 150 kg K₂O/ha assessed as statistically significant;
- With increasing fertilizers doses K:Mg ratio in the shoots and leaves was more favorable.

Acknowledgement

This paper was realized as a part of the project (TR 31063) Application of new genotypes and technological innovation in fruit and grape production financed by the Ministry of Education and Science of the Republic of Serbia within the framework of the technological project research for the period 2011-2014.

BIBLIOGRAPHY

1. **Conradie, W.J.** (1994): *Vineyard fertilization, Handbook of a workshop on vineyard fertilization, Nietvoorbij Stellenbosch, South Africa, 41pp.*
2. **Garcia, M., Daverede, C., Gallego, P., Toumi, M.** (1999): *Effect of various potassium-calcium ratios on cation nutrition of grape grown hydroponically, Journal of Plant Nutrition, Vol. 22(3), pp. 417-425.*
3. **Gluhic, D., Herak Cusic, M., Petek, M., Čoga, L., Slunjski, S., Sinčić, M.** (2009): *The content of Mg, K and Ca ions in vine leaf under foliar application of magnesium on calcareous soils, Agriculturae Conspectus Scientificus 2:81-84.*
4. **Hlušek, J., Richter, R., Ryan, P.** (2002): *Vyživa a hnojeni zahradnich plodin, I ed. Praha, Zemedelec, pp, 81.*
5. **Ličina V.** (2009). *Agrohemija. Zavod za udžbenike i nastavna sredstva. Beograd.*
6. **Ličina, V., Jakovljević, M.** (1997): *The K/Mg ratio in grapevine organs growt on diferent soil types. Zemljište i biljka. Beograd. Vol. 46(2). pp. 178-182.*
7. **McLean, E. O., Hartwig, R. C., Eckert, D. J., Triplett, G. B.** (1983): *Basic cations saturation ratios as a basis for fertilizing and liming agronomic crops: II. Field studies. Agron. J. 75, pp. 635-639.*
8. **Nakalamić A.** (1991). *Modifikovana dvokraka asimetrična kordunica. Jugoslovensko vinogradarstvo i vinarstvo. Beograd. Br. 4, pp. 7-10.*
9. **Nakalamić A., Marković N.** (2009). *Opšte vinogradarstvo. Poljoprivredni fakultet. Beograd.*
10. **Džamić R., Stevanović D., Jakovljević M.** (1996). *Praktikum iz agrohemije. Naučna knjiga. Beograd.*
11. **Zatloukalova, T., Lošak, J., Hlušek, P., Pavloušek, P., Sedlaček, M., Filipčik, R.** (2011): *The effect of soil and foliar applications of magnesium fertilizers on yields and quality of vine (Vitis vinifera, L.) Grapes, Acta universitatis agriculturae et silviculturae mendelianae brunensis, Vol. 59(3), pp. 221-226.*