

## IMPACT OF IRRIGATION REGIME AND APPLICATION OF KAOLIN ON THE STOMATAL CONDUCTANCE AND LEAF WATER POTENTIAL OF PEPPER AND TOMATO

**Marija Ćosić<sup>1\*</sup>, Ružica Stričević<sup>1</sup>, Nevenka Djurović<sup>1</sup>, Ljiljana Prokić<sup>1</sup>, Milena Marjanović<sup>1</sup>, Djordje Moravčević<sup>1</sup>**

<sup>1</sup>University of Belgrade, Faculty of Agriculture, Nemanjina 6, 11080 Zemun, Serbia

\*e-mail: [c.marija@agrif.bg.ac.rs](mailto:c.marija@agrif.bg.ac.rs), telephone +38111 4413243

**Keywords:** irrigation regime, kaolin, stomatal conductance, leaf water potential.

### ABSTRACT

The effect of different irrigation regimes and the application of kaolin on stomatal conductance and leaf water potential of pepper (*Capsicum annuum* L.) and tomato (*Lycopersicon esculentum* Mill.), is discussed in the paper. The experiment was set up by a completely randomized block system with three replications. The peppers were observed in three, and tomatoes in two treatments of irrigation regimes. The kaolin treatments of both crops were: a) control without kaolin (C) and 5% kaolin suspension (K).

The results of these studies show that the application of deficit irrigation treatments and kaolin treatment have no statistically significant effect on stomatal conductance and leaf water potential of pepper and tomato plants.

### INTRODUCTION

Agriculture is the largest consumer of fresh water, and with the application of irrigation, the amount comes up to 70% of the total available water (Fischer et al., 2007). Successful agricultural production, especially of vegetables and fruits, cannot be imagined without irrigation. Given that water is a limited resource, research should go in the direction of finding the different systems and measures to conserve water and achieve satisfyingly economically profitable yields.

The application of deficit irrigation and kaolin suspension might be some of the options for mitigating negative effects of drought caused by climate change and for saving water in agricultural production. The application of deficit irrigation aims to save irrigation water, increase water use efficiency and achieve optimal yields (Topcu et al., 2007). By applying the strategy of deficit irrigation, crops are systematically exposed to moderate levels of stress due to a lack of water for a certain period or during the entire vegetation, which results in lowering yields, but also to cost savings and increased efficiency of water use (English and Raja, 1996; Pereira et al., 2002). Basically, the method of deficit irrigation is to reduce the amount of irrigation water to such extent to cause the adaptive response of plants to drought, enabling them to increase the efficiency of water use and maintain yields, while increasing the quality of fruit (Savic, 2008). Before a decision is made on the application of deficit irrigation regimes, it is important to assess its impact on different cultures on the basis of many years of experimental research (Scholberg et al., 2000; Igbadun et al., 2008).

Application of kaolin leads to shadowing of the plant (aboveground part and fruits) which leads to a reduction in water consumption. Many researches highlight the impact of kaolin in reducing sunburns on the fruits, such as pomegranates, apples, walnuts, citrus fruits, and tomatoes (Cantore et al., 2009; Glenn, 2012; Weerakkody et al., 2010; Pace et al., 2007). Kaolin-based particle film technology (Pft) has been developed over the past 15 years as a multi-functional, environment-friendly material, that provides effective insect control, mitigates heat stress, and contributes to the production of high-quality fruit and

vegetables, as well as being suitable for organic farming (Glenn and Puterka, 2005). Boar et al. (2015) have followed the effect of kaolin on a variety of crops (tomatoes, peppers, zucchini, eggplant, beans and clementines). Their results indicate that the kaolin had the biggest impact on reducing stomatal conductance, which contributed to reduction in transpiration and improved the water regime of plants, reducing the net assimilation. Kaolin can be, in addition to its positive effects in controlling pests and reducing heat stress, successfully used as antitranspirant to mitigate the effects of drought stress, salinity and preserving water in arid regions such as the Mediterranean (Boari et al., 2015).

The aim of this study was to investigate the effects of different irrigation regimes and application of kaolin on stomatal conductance and leaf water potential of peppers and tomatoes.

## MATERIALES AND METHODS

The experiment in Stara Pazova was set up by a completely randomized block system with three replications. The peppers were observed in three, and tomatoes in two treatments of irrigation regimes. In both culture treatments of kaolin served as a control treatment (C) with no treatment with 5% kaolin suspension and treated with 5% suspension of kaolin (K).

The pepper irrigation regimes and kaolin treatment were: i) full irrigation without kaolin (FC) and full irrigation with kaolin (FK), ensuring 100% of crop evapotranspiration (ET<sub>c</sub>); ii) deficit irrigation without kaolin (R1C) and deficit irrigation with kaolin (R1K), at 80% of ET<sub>c</sub> and iii) deficit irrigation without kaolin (R2C) and deficit irrigation with kaolin (R2K) at 70% of ET<sub>c</sub>. Two tomato irrigation treatments and kaolin treatment were studied: i) full irrigation without kaolin (FC) and full irrigation with kaolin (FK), covering 100% of ET<sub>c</sub>, and ii) deficit irrigation without kaolin (DC) and deficit irrigation with kaolin (DK), at 50% of ET<sub>c</sub>.

The stand density of pepper cv. *Elephant Ear* (paprika) was 60,000 plants per hectare. The tomato was of the determinate type, *Rio Grande* cultivar, with stand density of 30,000 plants per hectare. In both treatments the soil in the paired rows, under the plants, was covered with black plastic mulch over the 60%.

The soil in the study area is of the carbonate chernozem type, developed in loess. Its morphological, hydrophysical and agrichemical properties are conducive to farming. The quality of this soil is founded upon its cross-sectional depth, mechanical composition with nearly equal proportions of sand, silt and clay fractions, good texture – clayey loam (USDA, 2006), and superior physical and chemical properties.

The climate is continental, with Central European and Mediterranean components. The annual precipitation average over the past 20 years is 637 mm (maximum 911 mm and minimum 352 mm). The average precipitation total during the growing season (from April to September) is 366 mm (maximum 663 mm and minimum 193 mm). Figure 1 shows average monthly precipitation totals and mean monthly air temperatures.

Climate data were adopted from the Republic Hydro-meteorological Service for the meteorological station Surčin (44°49' N lat; 20°17' E long, 96 m a.s.l) which is considered representative for the study area. The average air temperatures and precipitation totals in the study area during the study period are also shown in Table 1.




Fig.1. Mean monthly air temperatures and precipitation totals.

Table 1

**Monthly mean temperature ( $T_{avg}$ ) and precipitation during the growing cycles of pepper and tomato**

Month/ Year	2011	
	$T_{avg}$ (°C)	$\Sigma$ Precipitation (mm)
May	16.9	94.8
June	21.3	23.0
July	23.3	41.1
August	23.7	5.3
September	21.8	28.9
Average/Sum	21.4	193.1

Irrigation was done by the drop-by-drop method. Table 2 shows irrigation depths and frequencies on all treatments. The irrigation frequency depended on current climate conditions (amount and distribution of rainfall and ETc).

Table 2

**Seasonal irrigation amount (I, mm) and watering number (W, n.)**

Bell pepper (cv <i>Elephant Ear</i> )			Tomato (cv <i>Rio Grande</i> )		
Year	2011		Year	2011	
Treatment	I (mm)	W (n.)	Treatment	I (mm)	W (n.)
F	450	25	F	270	15
R1	378	21	D	162	9
R2	270	15			


A 5% kaolin suspension (Surround WP<sup>®</sup>, 95% purity) was applied from the time of flowering to the time of ripening. The suspension was prepared *in situ*, immediately prior to use, and was applied by means of 15L backpack sprayers. The amount of kaolin varied depending on the plant's phenophase. Kaolin was applied seven times because of heavy rainfall in July, which washed out the kaolin from the plant.

Five times during the growing season the conductivity of stoma was measured ( $g_s$  in  $\mu\text{mol m}^{-2}\text{s}^{-1}$ ) using porometers (AP4, Delta-T, Cambridge, U.K.) and leaf water potential ( $\Psi_l$  in MPa) in a pressurized chamber (Soil Moisture Equipment Corp., Santa Barbara, CA, USA). The measurements were carried out on July 7; July 18; July 26; August 1 and August 12.

The collected experimental data were processed by statistical methods using IBM SPSS V. 20 statistical software. The effects of the studied factors, the irrigation regimes and the kaolin application, as well as their interactions were assessed applying the variance method for the two-factorial, completely random block system experiment.


## RESULTS AND DISCUSSIONS

The analysis of the measured conductivity of stoma during the growing seasons of peppers and tomatoes shows certain variations. It can be noted that the watering regime has an impact on stomatal conductance – the higher the level of irrigation, the higher stomatal conductance. When it comes to kaolin, a clear conclusion cannot be given because sometimes stomatal conductance in both cultures was sometimes higher in the treatments with kaolin than in the treatments without kaolin and vice versa (Fig. 2). What we can safely conclude is that the stomatal conductance in the treatment of full irrigation, with and without the use of kaolin on peppers and tomatoes is higher compared to other treatments.



**Fig.2. Stomatal conductance of peppers and tomatoes on all treatments**

Figure 3 shows the results of measurements of leaf water potential of peppers and tomatoes. The treatment of deficit irrigation affected the leaf water potential. In the conditions of intensive irrigation leaf water potential was lower in both cultures. As with stoma conductivity, a clear conclusion about the impact of kaolin on leaf water potential cannot be made, because it was sometimes lower in the treatments with kaolin compared to treatments without kaolin and vice versa.



**Fig.3. Leaf water potential of peppers and tomatoes on all treatments**

Figure 4 shows the relations between leaf water potential and stomatal conductance of peppers and tomatoes. Certain correlation can be noted, i.e. the higher the stomatal conductance, the potential is lower. Also, it can be seen that in the treatment of deficit irrigation the stomatal conductance is lower and leaf water potential is higher.




Fig.4. Relations between leaf water potential and stomatal conductance of peppers and tomatoes on all treatments





Fig. 5. Average values of stomatal conductance of peppers and tomatoes during the research period

Figure 5 shows the average values of stomatal conductance of peppers and tomatoes during the research period. In this case, we can see the influence of the regime of irrigation on stomatal conductance of peppers and tomatoes – the more intense the irrigation is, the stomatal conductance is higher. When it comes kaolin, a clear conclusion cannot be given because the stomatal conductance is sometimes higher in the treatments with kaolin (R1K treatment of peppers and FK of tomatoes), and sometimes lower. Figure 6 shows the average values of leaf water potential of peppers and tomatoes in all tested treatments. In both cultures there are very small differences in LWP between treatments with different supply of water and treatments with and without the use of kaolin.



**Fig. 6. Average values of leaf water potential of peppers and tomatoes in all tested treatments during the research period**

Based on the analysis of variance of average values of stomatal conductance and leaf water potential of peppers and tomatoes (Tab. 3) it can be concluded that neither the irrigation regime, nor the application of kaolin have significantly affected the SC and LWP of peppers and tomatoes.

The average stomatal conductance of pepper varies from  $463.4 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  in RK2 to  $659.3 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  in FC treatment. Leaf water potential ranges from  $-0.98 \text{ MPa}$  in FK treatment to  $-1.17 \text{ MPa}$  in R2C treatment. Here it can be noted that kaolin influenced the lowering of the potential which is in accordance with the results obtained by Boari et al. (2015), noting that kaolin influences the improvement of the water regime of the plant.

The average stomatal conductance of tomatoes goes from  $627.3 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  in DK to  $753.7 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  in FK. Here it is observed that kaolin can help save water as antitranspirant, in deficit irrigation treatments with and without the application of kaolin the stomatal conductance is lower. Leaf water potential in all treatments of tomato is quite balanced, it is the lowest in FC treatment ( $-0.53 \text{ MPa}$ ) and the highest in DC treatment ( $-0.60 \text{ MPa}$ ).

Many studies have come to similar results, showing that water stress affects the decrease in stomatal conductance and the increase in leaf water potential. Campos et al.

(2014) have found that water stress significantly reduces stomatal conductance and increases the leaf water potential of pepper. The same observations were made by Romero et al. (2010) with grapevine. Cantor et al. (2016) also point out that the maximum stomatal conductance of tomatoes can be found in full irrigation treatments.

Application of kaolin impacts the reducing of conductivity of stoma and evaporation (Cantore et al., 2009; Boari et al., 2014), which improves the water regime of the plant (Le Grange et al., 2004; Lombardini et al., 2005; and Shellie King, 2013a, b). For well-irrigated beans, kaolin influenced the decrease in stomatal conductance and transpiration (Boari et al., 2015). These results are consistent with the results of Tworkoski et al, (2002) also on the beans and Ananda Coomaraswamy et al, (2000); Cantor et al. (2009) on tomatoes and tea. Boari et al. (2015) point out that kaolin has the greatest impact on reducing stomatal conductance, which contributed to reducing transpiration and improving the water regime of plants, reducing the net assimilation.

**Table 3**

***The influence of irrigation and kaolin on SC and LWP of peppers and tomatoes***

Pepper									
SC ( $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )					LWP (MPa)				
Treatments of IR		Treatments of IR and KA		Avg.	Treatments of IR		Treatments of IR and KA		Avg.
ns		ns		ns	ns		ns		ns
<b>FC</b>	659.3	<b>FK</b>	557.4	608.4	<b>FC</b>	-1.10	<b>FK</b>	-0.98	-1.04
<b>R1C</b>	525.5	<b>R1K</b>	598.5	562.0	<b>R1C</b>	-1.01	<b>R1K</b>	-1.04	-1.03
<b>R2C</b>	505.9	<b>R2K</b>	463.4	484.7	<b>R2C</b>	-1.17	<b>R2K</b>	-1.12	-1.14
<b>Avg.</b>	563.6	<b>Avg.</b>	539.8	551.7	<b>Avg.</b>	-1.09	<b>Avg.</b>	-1.05	-1.07
Tomato									
SC ( $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )					LWP (MPa)				
Treatments of IR		Treatments of IR and KA		Avg.	Treatments of IR		Treatments of IR and KA		Avg.
ns		ns		ns	ns		ns		ns
<b>FC</b>	692.0	<b>FK</b>	753.7	722.9	<b>FC</b>	-0.53	<b>FK</b>	-0.58	-0.59
<b>DC</b>	671.1	<b>DK</b>	627.3	649.2	<b>DC</b>	-0.60	<b>DK</b>	-0.58	-0.56
<b>Avg.</b>	681.5	<b>Avg.</b>	690.5	686.1	<b>Avg.</b>	-0.57	<b>Avg.</b>	-0.58	-0.57

SC – Stomatal conductance, LWP - Leaf Water Potential, IR – Irrigation Regime, KA - Kaolin Application, ns - not significant

**CONCLUSIONS**

These studies led to the conclusion that neither the use of irrigation regime, nor the application of kaolin has significantly affected the stomatal conductance and leaf water potential of tomatoes and peppers. Therefore, in conditions of climate change and drought, irrigation can be deficit and kaolin can be applied without significantly affecting very important physiological parameters such as SC and LWP which was the main objective of this research.

Although not statistically significant, there are some differences in the conductivity of stoma and leaf water potentials between treatments differently supplied with water and treatments with and without the use of kaolin. Reduction of irrigation reduces stomatal conductance and the leaf water potential increases. Application of the kaolin on treatments of deficit irrigation in both cultures has led to some reduction in leaf water potential, which confirms noted that kaolin does influence the improvement of the water regime of the plant.

## ACKNOWLEDGMENT

The paper was produced within the scope of the TR 37005 Project funded by the Serbian Ministry of Education and Science.

## BIBLIOGRAPHY

1. **Anandacoomaraswamy, A., De Costa, W.A.J.M., Shyamalie, H.W., Campbell, G.S.,** 2000 - *Factors controlling transpiration of mature field-grown tea and its relationship with yield. Agric. Forest Meteorol.* 103, 375–386.
2. **Boari, F., Cucci, G., Donadio, A., Schiattone, M.I., Cantore, V.,** 2014 -*Kaolin influences tomato response to salinity: physiological aspects. Acta Agric. Scand., Sect. B: Plant Soil Sci.,* <http://dx.doi.org/10.1080/09064710.2014.930509>.
3. **Boari, F., Donadio, A., Schiattone, M. I., Cantore, V.,** 2015 -*Particle film technology: A supplemental tool to save water. Agric. Wat. Manage.* 147, 154–162.
4. **Campos, H., Trejo, C., Peña-Valdivia, C. B., García-Nava, R., Conde-Martínez, F. V., & Cruz-Ortega, M. R.,** 2014 -*Stomatal and non-stomatal limitations of bell pepper (Capsicum annuum L.) plants under water stress and re-watering: Delayed restoration of photosynthesis during recovery. Environmental and Experimental Botany,*98, 56-64.
5. **Cantore, V., Pace, B., Albrizio, R.,** 2009 -*Kaolin-based particle film technology affects tomato physiology, yield and quality. Environ. Exper. Bot.* 66, 279–288.
6. **Cantore, V., Lechkar, O., Karabulut, E., Sellami, M. H., Albrizio, R., Boari, F., Todorovic, M.** 2016 -*Combined effect of deficit irrigation and strobilurin application on yield, fruit quality and water use efficiency of “cherry” tomato (Solanum lycopersicum L.). Agricultural Water Management,* 167, 53-61.
7. **English, M., Raja, S.N.,** 1996 -*Perspective on deficit irrigation. Agric. Water Manage.* 32, pp. 1–14.
8. **Fischer, G., Tubiello, F.N., van Velthuizen, H., Wiberg, D.A.,** 2007 - *Climate change impacts on irrigation water requirements: effects of mitigation, 1990–2080. Technological Forecasting and Social Change* 74, 1083–1107.
9. **Glenn, D.M., Puterka, G.J.,** 2005 -*Particle films: a new technology for agriculture. Hortic. Rev.* 31, 1–44.
10. **Glenn, D. M.,**2012 -*The mechanisms of plant stress mitigation by kaolin-based particle films and applications in horticultural and agricultural crops. Hort. Science,* 47 (6), 710-711.
11. **IBM Corp. Released** 2011 - *IBM SPSS Statistics for Windows, Version 20.0.* Armonk, NY: IBM Corp.
12. **Igbadun, H.E., Salim, B.A., Tarimo, A.K.P.R., Mahoo, H.F.,** 2008 - *Effects of deficit irrigation scheduling on yields and soil water balance of irrigated maize. Irrigation Sci.* 27, pp. 11–23.
13. **Le Grange, M., Wand, S.J.E., Theron, K.I.,** 2004 - *Effect of kaolin applications on apple fruit quality and gas exchange of apple leaves. ActaHortic.* 636, 545–550.
14. **Lombardini, L., Harris, M.K., Glenn, D.M.,** 2005 - *Effects of particle film application on leaf gas exchange, water relations, nut yield, and insect populations in mature pecan trees. HortScience* 40, 1376–1380.
15. **Romero, P., Fernández-Fernández, J. I., & Martínez-Cutillas, A.,** 2010 - *Physiological thresholds for efficient regulated deficit-irrigation management in wine grapes grown under semiarid conditions. American Journal of Enology and Viticulture* 61(3), 300-312.
16. **Pace, B., Boari, F., Cantore, V., Leo, L., Vanadia, S., De Palma, E., Phillips, N.,** 2007 - *Effect of particle film technology on temperature, yield and quality of processing tomato..Acta Hort.* 758, 287-293.



17. **Pereira, L.S., Oweis, T., Zairi, A.**, 2002 - *Irrigation management under water scarcity. Agricultural Water Management*, 57, pp. 175–206.
18. **Savic, S.**, 2008 - *The physiological mechanisms of tomato (*Lycopersicon esculentum* L.) productivity regulation in drought stress conditions. PhD thesis, Faculty of Agriculture, University of Belgrade, Serbia.*
19. **Shellie, K., King, B.A.**, 2013a - *Kaolin particle film and water deficit influence red winegrape color under high solar radiation in an arid climate. Am. J. Enol. Vitic.* 64, 214–222.
20. **Shellie, K., King, B.A.**, 2013b - *Kaolin-based foliar reflectant and water deficit influence Malbec leaf and berry temperature, pigments, and photosynthesis. Am. J. Enol. Vitic.* 64, 223–230.
21. **Scholberg, J., McNeal, B.L., Jones, J.W., Boote, K.J., Stanley, C.D., Obreza, T.A.**, 2000 - *Growth and canopy characteristics of field-grown tomato. Agron. J.* 92, pp. 152–159.
22. **Tworowski, T.J., Glenn, D.M., Puterka, G.J.**, 2002 - *Response of bean to application of hydrophobic mineral particles. Can. J. Plant Sci.* 82, 217–219
23. **Topcu, S., Kirda, C., Dasgan, Y., Kaman, H., Cetin, M., Yazici, A., Bacon, M.A.**, 2007 - *Yield response and N-fertiliser recovery of tomato grown under deficit irrigation. European Journal of Agronomy*, 26, pp. 64–70.
24. **USDA**, 2006 - *Keys to Soil Taxonomy, Tenth edition. Soil Survey Staff, United States Department of Agriculture NRCS.*
25. **Weerakkody, P., Jobling, J., Infante, M.M.V., Rogers, G.**, 2010 - *The effect of maturity, sunburn and the application of sunscreens on the internal and external qualities of pomegranate fruit grown in Australia. Sci. Hortic.* 124, 57–61.