

SOME ADDITIONAL NOTES ON THE STATUS OF NATURA 2000 HABITATS FROM OLTENIA (ROMANIA)

R DUŢOIU DANIEL^{1*}, TEF NESCU DRAGO -MIHAIL¹, NICOLAE ION¹

¹ University of Craiova, Biology Department, 15, Libert ii Street, 200583

*Corresponding author, e-mail: radutoiudaniel@yahoo.com

Keywords: habitats, Natura 2000, sites, Oltenia, Romania.

ABSTRACT

Based on the numerous field trips made in various Natura 2000 sites within the territory of Oltenia, there were collected data that were then processed in the laboratory. After analysing the data, there were found some differences between the field reality and what is mentioned in the standard sheet of the respective areas.

The data presented in this paper make reference to certain habitats from Oltenia, habitats that either cover a reduced surface (9420 Alpine Larix decidua and/or Pinus cembra forests; 3230 Alpine rivers and their ligneous vegetation with Myricaria germanica; 6210 Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia)) or are not encountered, although they are mentioned in the standard sheet of certain areas (40C0* Ponto-Sarmatian deciduous shrubs; 91Q0 Western Carpathian calcicolous Pinus sylvestris forests).*

INTRODUCTION

The intense study of habitats began after 2000, following the introduction of Natura 2000 network in Romania. Habitats of Romania (Doni & al. 2005) and Interpretation Manual of European Union habitats in Romania (Gafta & Mountford 2008) represented the most important books for any student who had to study Natura 2000 habitats in a particular area of the country.

Few experts achieved research studies on the habitats of Oltenia (Niculescu L. & al. 2015; R duţoiu, 2015; R duţoiu & al., 2015; Bule Dragomir 2014, 2015; R duţoiu & tef nescu, 2014, R duţoiu & Marinescu, 2014).

MATERIAL AND METHODS

Based on the data from the specialized literature, there were made numerous field trips to identify correctly the vegetation characterizing the studied habitats.

The characterization of the typical vegetation of each habitat was performed taking into account the national literature (Sanda V. & al. 1980, Sanda V. & al. 2001, Sanda V., 2002, Sanda & al. 2007).

The codes and the description of the habitat types correspond to NATURA 2000 and the Romanian system of classification (Doni & al., 2005; Doni & al., 2006, Gafta & Mountford, 2008).

RESULTS AND DISCUSSIONS

The presence on smaller surfaces than those specified in the standard sheets of the protected areas or the mention of certain uncharacteristic Natura 2000 habitats within different sites on the territory of Oltenia, required the specification of certain issues that reflect the reality on the ground.

Among the inventoried habitats, we mention:

6210* Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia)

The dry grasslands falling within this habitat have a poor representation in Oltenia. Within the areas where they were identified, they present a rich and varied flora composition (Fig. 1). They form a real mosaic of plant associations.

The vegetation of this habitat belongs to *Danthonio - Brachypodietumpinnati* Soo (1946) 1947. In the areas where such grasslands develop, they are used as hayfields and, then, in autumn, they are grazed. In those areas where irrational grazing is practiced, it was observed that *Danthoniaalpina* disappeared, its place being taken by *Festucarupicola*.



Fig. 1. Physiognomy of this habitat within Racovi a settlement (orig.)

The association prefers gentle slopes with northern or northwestern exposure. Along with *Danthoniaalpina*, there are constantly met the following species: *Festucarupicola*, *Brachypodiumpinnatum*, *Koeleriamacrantha*, *Brizamedia*, *Trifoliummontanum*, *Festucavalesiaca*, *Lotus corniculatus*, *Trifoliumcampestre*, *Genistellasagittalis*, *Chamaecytisusalbus*, *Dorycniumherbaceum*, *Polygala major*, *Salvia pratensis*, *Thymus glabrescens*, *Plantago media*, *Asperulacynanchica*, *Achilleamillefolium*, *Leucanthemum vulgare* (Fig. 2).



Fig. 2. Aspect of the phytocoenoses of this association within Cerna Sat settlement (orig.)

3230 Alpine rivers and their ligneous vegetation with *Myricariagermanica*

It is a habitat identified along certain rivers in Oltenia. It develops on alluvial soils with a high level of moisture and terrains uncovered by floods. In some areas, it has a poor representation (ROSCI0129 North of Western Gorj, ROSCI0128 North of Eastern Gorj),

while in others, where it has a continuous development, it is not mentioned (ROSCI0239 - Tarnovu Mare - Latori a).

The increased heterogeneity and low constancy of the species included in the composition of this habitat are mainly due to the repeated floods that occur during the growing season, by which new species are brought, while others are eliminated.

Due to the root system of the plants characteristic to this habitat, which is very well developed, the vegetation of these places plays a major role in stabilizing the alluvia, thereby contributing to the vegetation succession on the riverbank.

The vegetation characteristic to this habitat belongs to the association *Salicipurpureae - Myricarietum* Moor 1958 (Gafta& Mountford 2008).

The areas covered by phytocoenoses edified by *Salix purpurea* and *Myricariagermanica* are located from the sub-Carpathian hills to the middle of the boreal level of the sites North of Western Gorj, North of Eastern Gorj and Târnovu Mare - Latorița. They prefer the gravel areas of the floodplains of mountain valleys. As well as all the phytocoenoses that prefer these places, they manifest a strong pioneering character. The modification of the floristic composition is induced by the numerous floods that occur during the year.

In the area Târnovu Mare –Latori a, this habitat had a very good representation in the years 2013-2014, but, in the summer of 2014, because of flash floods, the area of this habitat reduced considerably. This phenomenon was observed both along the valley of the Repede Stream and that of the Latori a River.

Salix purpurea has a better representation at lower altitudes and *Myricariagermanica* within beech forests (Fig. 3).



Fig. 3. Detail of a specimen of *Myricariagermanica* on the Valley of the Repede Stream (orig.)

91Q0 Western Carpathian calcicolous *Pinussylvestris* forests

This habitat is edified by one of the oldest tree species in our flora. Tens of thousands of years ago, in the forests of Romania and Central Europe, the dominant species was the Scots pine (*Pinussylvestris*). This is proven by pollen analyses conducted by specialists over time.

In Romania, as well as within the territory investigated by us, this habitat has little representation. Whether at national level these relict habitats are found only in Trascău Mountains, the Bicăz Gorges, Vrancea, Leaota and Cozia Mountains, within the site ROSCI0128 North of Eastern Gorj, they are located in the eastern part of the area, on very

small surfaces (eg. TheCernaOlte ului Valley, next toStâni oara Peak, on the slope withwestern exposure).

The lack of valuable specimens in these areas enables us to say that, in these forests, there were made either cuttings and, in this case, present forests are remnants of the old ones, or are plantations made in the 80s, plantations that grow well in the climatic conditions registered in the region.

The absence from the floristic composition of the species that give the name of the two associations (*Sesleriorigidae* - *Pinetumsylvestris*Cs röset al. 1988 and*Daphnoblageyanae* - *Pinetumsylvestris*ColdeaetPop 1988) that characterize the vegetation of this habitat in the entire country (Gafta& Mountford, 2008), make us mention that this habitat cover a limited surface in Oltenia. Besides the classical sites mentioned in specialized literature in this part of the country, this habitat covers only small emerging surfaces. The analysis of the floristic composition enables us to say that, within the sites from Oltenia (except for the surfaces located within Cozia Massif) this habitat is an artificial habitat, resulting from the plantation of the dominant species.

9420 Alpine *Larix decidua* and/or *Pinus cembra* forests

In the Alps, this habitat is characteristic tothe subalpine zone and rarely develops in the zone of spruce-fir forests. In most cases,*Larix decidua* or *Pinus cembra*are the dominant species in these forests. The two species can form pure or mixed forests and can be associated with *Picea abies*.

In Romania, this habitat is rare, being concentrated in five relict centres, some of them being located at much lower altitude. It is about Ceahl u, Ciuca , Bucegi, Latori a and Trasc u (the northern part–Vidolm and Piatra Secuiului) Mountains, respectivelyMare (Big) Mountain (Sc ri a-Belioara) (F rca & al., 2013,Merce 2013).

All natural larch forests from our country are conditioned by the occurrence of limestone or calcareous conglomerates.

Besides larch, in the aforementioned areas, there also appear spruce, silver fir, beech, birch, mountainmaple.

In all locations, the herbaceous layer is characterized by the presence of *Saxifragacuneifolia* subsp. *robusta* (Fig. 4) together with an interesting mix of species characteristic to calcareous grassland, as well as acidophilus species such as the Carpathian *Sesleriarigida*.



Fig. 4. *Saxifragacuneifolia* subsp. *robusta* from ROSCI0239 (orig.)

In Romania, *Larix decidua* forests cover about 30,000 hectares, most of them being plantations grown from seeds imported from Austria (St nescu, 1979). Only about 4,500 ha are considered natural.

The larch forest from the site includes a natural protected area of national interest corresponding IUCN category IV. It is on the right side of the Latori a River, at an altitude between 800 and 1,350 m. This is one of 6 natural larch locations in Romania, as mentioned by Gh. Ploaie (2012).

These forests also resulted from plantations made decades ago when clearcutting was practiced in the area. In some bibliographic sources, it is also mentioned *Pinus cembra* in these areas, but we did not identify it. Perhaps it grew in the area before the clearcutting made in the 80s (data reported orally by the forester who watches these forests). It usually grows on strongly inclined western slopes.

40C0* Ponto-Sarmatian deciduous shrubs

These are shrubs that develop well in dry areas. Their extension and coenotic-structural diversification is determined by soil and strongly influenced by human activity.

In "Oltenia Forest Steppe", this habitat is mentioned, but it was not identified in the field, either here or in the western part of the country. The shrubs of this site have a different vegetation compared to the areas in Dobrogea, where there is also found this habitat, but their landscape is typical to the steppe. There, the vegetation is edified by the associations *Asphodelinoluteae-Paliuretum* and *Rhamnocatharticae-Jasminietum fruticantis*, associations that are not present within Oltenia.

In all Natura 2000 sites where this habitat is mentioned, it has to be specified that Oltenia shrubs belong to the following categories: R3122 Ponto-Pannonic blackthorn (*Prunus spinosa*) shrubs and hawthorn (*Crataegus monogyna*) (40A0 *), Emerald priority habitat.

CONCLUSIONS

The habitats analysed in the present paper belong to all types of vegetation in Romania: arboreal (9420 Alpine *Larix decidua* and/or *Pinus cembra* forests in the mountainous region, 91Q0 Carpathian relict calcicolous *Pinus sylvestris* forests), shrub (3230 Alpine rivers and their ligneous vegetation with *Myricaria germanica*, 40C0 * Ponto-Sarmatian deciduous shrubs) and herbaceous (6210 * Semi-natural dry grasslands and scrubland facies on calcareous substrates (*Festuco-Brometalia*)).

9420 and 91Q0 habitats are rare or their presence is uncertain in certain areas of Oltenia; 3230 covers small surfaces in various protected areas, surfaces that have a different physiognomy conditioned by the places where they vegetate and by the climatic conditions during the year; 40C0* does not cover any surface within the territory of Oltenia considering the characteristic vegetation of this habitat, while in case of the habitat 6210*, taking into account its reduced surfaces in various Natura 2000 sites on the territory of Oltenia, there are required urgent measures to facilitate the expansion of these grasslands.

BIBLIOGRAPHY

1. **Bu e Dragomir Luminita, Niculescu Mariana.** 2015. *Physiological particularities of hygrophilous plants from Domogled-Cerna Valley National Park* Analele Universit ii din Craiova, seria Agricultur – Montanologie – Cadastru Vol. XLV (1): 51-56, <http://anale.agro-craiova.ro/index.php/aamc/index>
2. **Bu e Dragomir Luminita, Nicolae Ion.** 2014. *Ecophysiological aspects regarding the „Sibleac,, associations from Iron Gates Natural Park*, Analele Universit ii din Craiova, seria Agricultur – Montanologie – Cadastru Vol. 44, <http://anale.agro-craiova.ro/index.php/aamc/article/viewFile/114/108>

3. **Doni , N., Popescu, A., Pauc -Com nescu, M., Mih ilescu, S., Biri I.-A.** 2005. *Habitatale din România*, Edit. Tehnic Silvic Bucure ti. 496 pag.
4. **Doni , N., Popescu, A., Pauc -Com nescu, M., Mih ilescu, S., Biri I.A.** 2006. *Habitatalele din România*. Modific ri conform amendamentelor propuse de România i Bulgaria la Directiva Habitatale (92/43/EEC), Edit. Tehnic Silvic , Bucure ti.
5. **Gafta, D., Mountford, O.** (coord.), 2008. *Manual de interpretare a habitatelor Natura 2000 din România*, Edit. Risoprint, Cluj-Napoca.
6. **F rca Sorina, Tan u I. & Turtureanu P.D.** 2013. *Larix decidua Mill. in Romania: current and past distribution, coenotic references, and conservation status*. Contrib. Bot. XLVIII: 39-50.
7. **Merce O.** 2013. *Specific management measures for the forest habitats 9410 and 9420*. Journal of Horticulture, Forestry and Biotechnology, Vol. 17(2): 264- 267.
8. **Niculescu L., Niculescu M., Soare R. & Bonea D.** 2015. *Plant communities characteristic for the mountainous and subalpine springs and streams in the Parâng Mountains (Southern Carpathians), Romania*. Journal of Environmental Protection and Ecology 16, no. 4: 1364-1370.
9. **Ploaie Gh.** 2012. *Mun ii C p âinii. Ghidturistic*. Edit. Conphys, Rm. Vâlcea, 246 pag.
10. **R duțoiu D. & tef nescu M.** 2014. *Site of vascular flora Târnovu Mare – Latori a Vâlcea -Romania*. Annals of the University of Craiova. Seria Biologie, Horticultur , Tehnologia prelucr rii produselor agricole, Ingineria Mediului. Vol. XIX (LV): 563-570.
11. **R duțoiu D. & Marinescu E.** 2014. *Study of hygrophilous vegetation from Târnovu Mare-Latori a, Vâlcea - Romania*. Annals of the University of Craiova. Seria Biologie, Horticultur , Tehnologia prelucr rii produselor agricole, Ingineria Mediului. Vol. XIX (LV): 571-576.
12. **R duțoiu D.** 2015. *The conservation status of grassland habitats belonging to protected areas from Oltenia 'Natura 2000' site, Romania*. Not. Sci. Biol. 7 (4): 430-434.
13. **R duțoiu D., tef nescu M. & P unescu C.** 2015. *About the aquatic and paludous habitats from Oltenia (Romania)*. Annals of the University of Craiova. Seria Biologie, Horticultur , Tehnologia prelucr rii produselor agricole, Ingineria Mediului. Vol. XX (LVI): 589-594.
14. **Sanda V. & al.** 1980. *Cenotaxonomia i corologia grup rilor vegetale din România*. Stud. Com. (24) Supliment tiin ele Naturii Muzeul Brukenthal: 171 pag. Sibiu.
15. **Sanda V., Popescu A. & Stancu Daniela Ileana.** 2001. *Structura cenotic i caracterizarea ecologic a fitocenozelor din România*. 359 pag. Edit. CONPHIS, Bucure ti.
16. **Sanda V.** 2002. *Vademecum ceno structural privind covorul vegetal din România*. 331 pag. Edit. Vergiliu. Bucure ti.
17. **Sanda V., R du oi u D., Baraba N., Claudia Bi -Nicolae, Irina Blaj-Irimia.** 2007. *Breviar fitocenologic (partea III-a)*. Edit. Sitech. Craiova. 286 pag.
18. **Sanda V., R du oi u D., Burescu P., Irina Blaj-Irimia.** 2007. *Breviar fitocenologic (partea IV-a)*. Edit. Sitech. Craiova. 244 pag.
19. **St nescu V.** 1979. *Dendrologie*. Edit. Did. Ped., Bucure ti.