

BEHAVIOR OF SOME POTATO VARIETIES UNDER THE INFLUENCE OF THE THERMO-HYDRIC STRESS FACTORS SPECIFIC TO THE SANDY SOILS AREA

Diaconu Aurelia¹, Ploae Marieta¹, Paraschiv Alina¹

¹Research - Development Center for Field Crops on Sandy Soils, Dabuleni, Dolj County, Romania, Phone: +40251334402, Fax: +40251334347, E-mail ccdcpndabuleni@yahoo.com

Keywords: stress, potato, photosynthesis, foliar transpiration, sands

ABSTRACT

During the vegetation period, was analyzed the diurnal and seasonal variation of the physiological processes (the rate of photosynthesis and foliar transpiration) at 4 varieties and 2 lines of potato grown on sandy soils. The rate of photosynthesis showed a diurnal variation, being influenced by the active radiation in photosynthesis and air and soil humidity, as well as by the studied varieties and lines. In the tuber growth phase, the average of photosynthesis oscillated between 12,49 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at the Castrum variety and 21,03 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at the L 15-1677/31. At this stage of vegetation, the L 15 line was net detached of the other varieties, having a maximum photosynthesis which has positively influenced the production of tubers, recorded at 45 days after the emergence of plants (42,4 tons/ha). The high air temperature of 37,3 °C and the relative air humidity below 30% increased the evaporation of water through foliar transpiration at all varieties and lines studied. High values at the transpiration rate were recorded at 12-15 o'clock, when the action of stress factors was maximal, at the Sarmis variety (8,35 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$), and at line L 15-1677/31 (8,74 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$).

INTRODUCTION

Researches on the behavior of potato varieties and the influence of agro-technical factors (irrigation and fertilization) were performed in other potato culture areas by researchers: Chichea and Săninioiu (1994), Berindei and Chichea (1997), Draica (1995). Research by Carpentieri 1990 showed that photosynthetic activity remained stable up to 30 °C then decreased and inhibited to 40 °C. Research on Hammes potato in 1990 showed that raising the temperature from 15 °C to 40 °C resulted in a decrease in the net intensity of the photosynthesis process by 37 % compared to that determined at 20 °C. Under conditions of thermal stress the photosynthesis process is inhibited by decrease in Ribuloso-diphosphate-carboxylase enzyme activity after Sage and Cubien in 2007. Drought determines the concentration of the soil solution, reduces foliar absorption by accumulation of abscisic acid that closes stomata and increases 100 times in water stress (Salisbury and Ross-1991). Active photosynthetic radiation plays a role in the induction of photosynthesis as well as foliar transpiration, by the photoactive actions of opening the stomata and by increasing the temperature of the leaves (Burzo I. 2004). Atmospheric and pedological drought limits the metabolism of plants that enter into thermo-hydric stress (Petcu Elena 2008).

MATERIALS AND METHODS

During the vegetation period of 2017 year, were studied 4 varieties and 2 potato lines , with determinations carried out on vegetation phases and in 3 moments during the day, regarding:

- the active radiation in photosynthesis;
- the rate of photosynthesis;
- air temperature at leaves level;
- the foliar transpiration rate.

All this determinations where made with the LC PRO + portable device.

RESEARCH RESULTS

During the vegetation period at 45 days after the emergence of potato plants (Table 1) PAR (Photosynthetic Active Radiation) was oscillating at 9 o'clock between 1315 and 1328 $\mu\text{mol}/\text{m}^2/\text{s}$, at 12 o'clock between 1823 and 1860 $\mu\text{mol}/\text{m}^2/\text{s}$ and between 1454-1864 $\mu\text{mol}/\text{m}^2/\text{s}$ at 15 o'clock. The L14, L15 lines and Castrum variety recorded peak values of the active radiation in photosynthesis at the time of the determinations.

The rate of photosynthesis (Table 1) was influenced by the high air temperature of 37,2 °C and low atmospheric humidity to 30 %.

The rate of photosynthesis showed a diurnal variation under the influence of climatic factors in the area.

Photosynthesis rate values oscillated at 7 o'clock between 7,30 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at Castrum variety and 23,82 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at L 15-1677/31 line; at 12 o'clock between 15,07 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at Marvis variety and 28,02 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at the Braşovia variety, and at 15 o'clock between 7,37 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at Marvis variety and 18,09 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at the Sarmis variety.

The Braşovia variety and the L15 line recorded maximum values at the rate of photosynthesis at this stage of vegetation. The daily average values oscillated between 12,49 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at the Castrum variety and 21,03 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at line L 15 - 1677/31. The L15 line showed maximum value at the rate of photosynthesis, being double compared to the Marvis and Castrum varieties.

Table 1

Diurnal variation of photosynthesis rate at potato varieties at 45 days after the emergence

Variety	PAR $\mu\text{mol}/\text{m}^2/\text{s}$	Photosynthesis rate $\mu\text{mol CO}_2/\text{m}^2/\text{s}$	PAR $\mu\text{mol}/\text{m}^2/\text{s}$	Photosynthesis rate $\mu\text{mol CO}_2/\text{m}^2/\text{s}$	PAR $\mu\text{mol}/\text{m}^2/\text{s}$	Photosynthesis rate $\mu\text{mol CO}_2/\text{m}^2/\text{s}$	Daily average of photosynthesis $\mu\text{mol CO}_2/\text{m}^2/\text{s}$
	9 o'clock		12 o'clock		15 o'clock		
Marvis	1325	19.49	1833	15.07	1454	7.37	13.97
Braşovia	1317	20.69	1823	28.02	1500	13.35	20.68
Castrum	1315	7.30	1860	21.77	1797	8.42	12.49
Sarmis	1312	19.40	1830	22.55	1600	18.09	20.01
L 14-1574/4	1322	17.84	1833	18.26	1864	9.56	15.22
L 15-1677/31	1328	23.82	1840	20.98	1506	18.31	21.03

The foliar transpiration rate (Table 2) was influenced by climatic factors (air temperature, air and soil humidity) and varieties and lines studied. The daily variation in foliar transpiration rate recorded the following values:

- at 9 o'clock the values varied between 4,99 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ at Castrum variety and 2,64 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ at line L15-1677/31;
- at 12 o'clock the values of the transpiration rate oscillated between 8,40 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ at the Sarmis variety and 10,14 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ at line L 15-1677/31;
- at 15 o'clock values fluctuated between 4,56 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ at the Marvis variety and 10,72 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ at line L 15-1677/31.

Line L 15-1677/31 recorded maximum values at the foliar transpiration rate at all times of the determinations and made very good use of the evaporated water by foliar transpiration because the recorded photosynthesis was maximum under the same climatic conditions. The daily average value recorded a minimum of 6,08 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ in the Marvis variety and a maximum of 8,85 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$ at line L 15-1677/31.

In this vegetation phase, the L15 line is net detached of the other varieties, having maximum photosynthesis which positively influenced the tubers production at 45 days after emergence and recorded 42,42 tons / ha.

Table 2

Diurnal variation of leaf transpiration rate at potato varieties at 45 days after the emergence

Variety	Air temperature at leaves level (°C)	Transpiration rate mmoli H ₂ O/m ² /s	Air temperature at leaves level (°C)	Transpiration rate mmoli H ₂ O/m ² /s	Air temperature at leaves level (°C)	Transpiration rate mmoli H ₂ O/m ² /s	Daily average of leaf transpiration
	9 o'clock		12 o'clock		15 o'clock		
Marvis	30.9	5.00	36.0	8.69	37.2	4.56	6.08
Brașovia	31.0	5.12	34.9	9.15	36.7	9.22	7.83
Castrum	30.7	5.40	35.0	9.01	37.8	7.83	7.41
Sarmis	29.6	4.99	34.5	8.40	37.6	8.79	7.39
L 14-1574/4	31.5	5.34	36.6	8.97	37.2	6.66	6.99
L 15-1677/31	31,8	5.69	36.6	10.14	37.3	10.72	8.85

Figure 1 shows the variation of photosynthesis and foliar transpiration rate in relation to air temperature and studied varieties. Under the same climatic conditions, the Brașovia, Sarmis and L 15-1677/31 varieties showed higher values at the rate of photosynthesis. The foliar transpiration rate is more influenced by the air temperature, of 34 – 35 °C.




Fig.1 Daily average of photosynthesis and foliar transpiration depending on temperature and variety at 45 days after the emergence

Tuber production at the first harvest ranged between 13,56 tonnes/ha at the Castrum variety and 44,34 tonnes/ha on line L15-1677/31.

Figure 2 shows the translocation of assimilates, which occurred either to vegetative growth or to tuber production. At the two lines the assimilates are found in tuber production, compared to Castrum where the assimilates are directed to vegetative growths, at the expense of production.




Fig. 2 Translocation of assimilates into the vegetativ growths and tubers at 45 days after emergence

In the intensive growth phase of the tubers at 55 days from emergence (Table 3) PAR oscillated between 1240 μmol/m²/s at Marvis variety and 1263 μmol/m²/s at Castrum variety at 9 o'clock, between 1570 μmol/m²/s at Sarmis variety and 1704 μmol/m²/s at the Brașovia variety at 12 o'clock and between 1290 μmol/m²/s at the Marvis variety and 1770 μmol/m²/s at the Sarmis variety at 15 o'clock.

The photosynthesis rate values oscillated:

- at 9 am, between 13,61 μmol CO₂/m²/s at Marvis variety and 27,04 μmol CO₂/m²/s at line L 15-1677/31.

- at 12 o'clock between 6,39 μmol CO₂/m²/s at line L 14-1574/4 and 17,01 μmol CO₂/m²/s at the Sarmis variety.

- at 15 o'clock between 1,15 μmol CO₂/m²/s at line L 14-1574/4 and 18,38 μmol CO₂/m²/s at L 15-1677/31.

The average daily value oscillated between 7,71 μmol CO₂/m²/s at L 14-1574/4 and 20,61 μmol CO₂/m²/s at L 15-1677/31.

In this phase of vegetation the Sarmis variety and line L 15-1677/31 recorded high values at the rate of photosynthesis compared to other varieties.

Table 3

Diurnal variation of photosynthesis rate at potato varieties at 55 days after the emergence (12.06.2017)

Variety	PAR μmol/m ² /s	Photosynthesis rate μmol CO ₂ /m ² /s	PAR μmol/m ² /s	Photosynthesis rate μmol CO ₂ /m ² /s	PAR μmol/m ² /s	Photosynthesis rate μmol CO ₂ /m ² /s	Daily average of photosynthesis
	9 o'clock		12 o'clock		15 o'clock		
Marvis	1240	13.61	1689	8.95	1290	4.01	8.85
Brașovia	1261	16.86	1704	7.26	1779	3.20	9.10
Castrum	1263	22.70	1648	15.86	1779	9.15	15.90
Sarmis	1253	15.30	1570	17.01	1884	12.13	14.81
L 14-1574/4	1245	15.59	1683	6.39	1358	1.15	7.71
L 15-1677/31	1255	27.04	1658	16.43	1740	18.38	20.61

In the intense growth phase of the tubers, the air temperature oscillated between 30 and 37,3 °C, influencing the loss of water by evaporation at the leaves level.

The diurnal variation of leaf transpiration rate (Table 4) recorded the following values:

- at 9 o'clock values ranged between 3,07 mmol H₂O/m²/s at the Marvis variety and 5,11 mmol H₂O/m²/s at Castrum variety;

- at 12 o'clock the values of the transpiration rate oscillated between 4,43 mmol H₂O/m²/s at Brasovia and 8,35 mmol H₂O/m²/s at the Sarmis variety;

- at 15 o'clock values fluctuated between 2,06 mmol H₂O/m²/s at line L14 and 8,74 mmol H₂O/m²/s at line L15-1677/31.

The maximum transpiration rate was recorded at 12-15 o'clock, at the Sarmis variety and at line L 15-1677/31, and the minimum was recorded at 9 o'clock at L 14 - 1574/4.

The daily average value recorded a minimum of 3,37 mmol H₂O/m²/s at L14-1574/4 and a maximum of 6,64 mmol H₂O/m²/s at L15-1677/31.

The high air temperature of 37,3 °C and the air humidity below 30 % increased the evaporation of water through foliar transpiration in all varieties and lines studied. The L 15-1677/31 line efficiently used evaporated water, recording high values at the rate of photosynthesis.

Table 4

Diurnal variation of leaf transpiration rate at potato varieties at 55 days after the emergence

Variety	Air temperature at leaves level (°C)	Transpiration rate mmoli H ₂ O/m ² /s	Air temperature at leaves level (°C)	Transpiration rate mmoli H ₂ O/m ² /s	Air temperature at leaves level (°C)	Transpiration rate mmoli H ₂ O/m ² /s	Daily average of leaf transpiration
	9 o'clock		12 o'clock		15 o'clock		
Marvis	30.2	3.07	35.6	7.02	36.4	3.74	4.61
Brașovia	30.4	4.31	37.0	4.43	35.5	2.30	3.68
Castrum	30.4	5.11	37.2	6.80	36.1	4.12	5.34
Sarmis	30.0	3.68	36.7	8.35	36.5	6.63	6.28
L 14-1574/4	30.7	3.32	35.3	4.75	37.3	2.06	3.37
L 15-1677/31	30.8	4.98	35.0	6.21	37.3	8.74	6.64

Analyzing the results obtained at 55 days after the emergence, it was observed that at a relatively constant and quite high average temperature, the photosynthesis values oscillated quite a lot, depending on the variety, detaching the Sarmis, Castrum and L 15-1677/31. Lower values recorded the Marvis, Brasovia and L14-1574/4 varieties, which demonstrate the influence of the variety on physiological processes.

Tuber production at second harvest oscillated between 20,94 tonnes/ha at Marvis variety and 45,62 tonnes/ha at line L15-1677/31 (Table 5).

Table 5

Photosynthetic yield at second harvest (55 days after the emergence)

Variety	Photosynthesis (μmol CO ₂ /m ² /s)	Weight of vegetative part of plants (t/ha)	Tuber production (t/ha)
Marvis	8,85	7.50	20.94
Brașovia	9,10	13.44	34.02
Castrum	15,90	21.72	34.62
Sarmis	14,81	16.20	37.74
L14 – 1574/4	7,71	14.16	37.32
L15 – 1677/31	20,61	14.88	45.62

CONCLUSIONS

Climatic factors during the vegetation period (air temperature above 35 °C, air humidity below 30 %) influenced the speed and pace of the physiological processes under study.

The rate of photosynthesis showed a diurnal variation, being influenced by the active radiation in photosynthesis and air and soil humidity, as well as by the studied varieties and lines.

In the tuber growth phase at 45 days after emergence, the mean daily value oscillated between 12,49 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at the Castrum variety and 21,03 $\mu\text{mol CO}_2/\text{m}^2/\text{s}$ at line L 15 – 1677/31. The L15 line showed maximum value at the rate of photosynthesis, being double compared to the Marvis and Castrum varieties. In this vegetation phase, the L15 line is net detached of the other varieties, having maximum photosynthesis which positively influenced the tubers production at 45 days after emergence and recorded 42,42 tons/ha.

Line L 15-1677/31 recorded maximum values at the foliar transpiration rate at all times of the determinations and very efficiently used the evaporated water through foliar transpiration because the recorded photosynthesis was maximum under the same climatic conditions.

High values at the transpiration rate were recorded at 12-15 pm, when the action of stress factors was maximal, at the Sarmis variety (8,35 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$) and at line L 15-1677/31 (8,74 $\text{mmol H}_2\text{O}/\text{m}^2/\text{s}$)

Drought-resistant and high-yielding potato lines will be expanded into crops on sandy soils.

BIBLIOGRAPHY

1. **Berindei M., Chichea I.**, 1997. *Cartoful timpuriu*. Edit. Grand Bucuresti pg.18-23
2. **Carpentieri,R.**,1999, *Effect of high-temperature Stresson the photosyntetic Apparatus*. Edit.Inc.New York, Basel.
3. **Chichea I., Săninoiu M.**, 1994. *Influența epocii de plantare asupra producției la cartoful timpuriu cultivat pe 2 tipuri de sol*. Edit. Alma, Craiova.
4. **Chichea I.**, 2000. *Cartoful timpuriu si de vara*. Edit. Alma Craiova.
5. **Hammes,P.S.** 1990 -*Net photosynthetic rate of potato hgh temperature*. Journal potato Research,vol.33, 515-520.
6. **Petcu Elena**, 2008, *Impactul schimbărilor climatice asupra plantelor: SECETA. DOMINO*.
7. **Sage and Kubien**, 2007. *The temperature response of C3 and C4 photosynthesis*.Enviroment 30,1086-1107.
- 8.**Salisbury,F.B.Ross**, 1991. *Plant Physiology*. Belmont, California.