

ANTIOXIDANT ENZYMES ACTIVITIES AND PROLINE CONTENT IN LEAVES OF SALIX SPECIES GROWN ON FLY ASH DUMPS

BABEANU CRISTINA¹, DODOCIOIU ANA MARIA^{2*}

1 University of Craiova, Faculty of Science, 107i Calea Bucuresti, Craiova, Romania

2 University of Craiova, Faculty of Horticulture, 13 A.I.Cuza Street, Craiova, Romania

*Corresponding author e-mail: anadodocioiu@gmail.com

Keywords: *Salix*, catalase, peroxidase, proline, fly ash

ABSTRACT

For remediation and re-cultivation of polluted soils, including fly ash dumps, stress-tolerant plants are needed, one of the species used for this purpose being the willow tree. A first response of the plant to different types of stress is the activation of the enzymatic and non-enzymatic antioxidant system for the neutralization of excess reactive oxygen species.

The purpose of this paper is to evaluate the activity of catalase and peroxidase and proline content in leaves of 14 clones of Salix (7 Romanian and 7 Swedish) grown on fly ash dumps from Isalnita. The obtained data were compared to those determined for control cultures from Radovan. The samples were collected in July and were determined the catalase and peroxidase activity by colorimetric method. The proline content was determined from sulfosalicylic acid extract by colorimetric method with ninhidric acid as reagent using L proline as standard.

The results obtained show an increase in peroxidase activity and in proline content (with few exceptions) in plants grown on ash dumps in comparison with the control plantation. The increase in antioxidant enzymes activities suggests a state of oxidative stress, the plants activating a defensive system. Oxidoreductase activity and proline content might be used as biomarkers of tolerance/adaptation of Salix species on degraded soils.

INTRODUCTION

Due to the global energy crisis, renewable energy resources, represented by crop biomass of SRC (Short rotation coppice) are a sustainable option. One of the species used for this purpose is the willow and a particular interest for the energy willow culture is observed (Hernea et al., 2015, Hernea et al., 2016). This interest in obtaining biomass can be connected with the remediation of polluted soils, including fly ash dumps (Soare et al., 2017).

Fly ash characteristic depends on physicochemical properties of coal, coal combustion process storage and climate (Ibrahim, 2015). Numerous scientific studies report that ash deposits contain potentially toxic elements like heavy metals in quantities that for some of them exceed the alert limit (Popa et al., 2013, Soare et al., 2010). These heavy metals are known to induce a state of oxidative stress in plants limiting their survival and growth.

A first response of the plant to this type of stress is the activation of the enzymatic and non-enzymatic antioxidant system for the neutralization of excess reactive oxygen species. If they are not neutralized, reactive oxygen species can cause lipid peroxidation,

membrane injury, protein degradation, enzyme inactivation, pigment bleaching and disruption of DNA strands (Singh and Tuteja, 2010).

The purpose of this paper is to evaluate the activity of some antioxidant enzymes (catalase and peroxidase) and proline content in leaves of 14 clones of *Salix* (7 Romanian and 7 Swedish) grown on fly ash dumps from Isalnita.

MATERIALS AND METHODS

The biological material was represented by ten *Salix* clones, seven Romanian: RO892, RO1077, RO1082, Cozia1, Fragisal, Pesred, Robisal and seven Swedish clones: Inger, Tordis, Jorr, Olof, Tora, Torhild and Svem. The behavior of investigated clones has also been studied in other areas in Romania (Babeanu et al., 2017, Hernea et al., 2016, Soare et al., 2015).

The studied plantations are located on fly ash deposits (N 44°24'41" E 23°41'07"), resulting from coal burning to Isalnita Power Station, near Craiova, Romania. These deposits are characterized by low fertility and many rehabilitation eco-technologies have been experimented (Babeanu et al., 2004; Popa et al., 2010).

Enzyme assays: Fresh tissue was homogenated with 0.1 M phosphate buffer, pH 7.0 (1:20 w:v) containing 0.1 mM EDTA. Homogenates were centrifuged for 20 min at 10,000 r.p.m. and the supernatants were used for enzyme assay.

Total soluble peroxidase activity (guaiacol-type E.C.1.11.1.7) was assayed by measuring the increase in A_{436} due to the guaiacol oxidation and their activity was expressed as $\Delta A/\text{min}/1\text{g}$ fresh weight (Babeanu et al., 2010).

Catalase activity (E.C.1.11.1.6) was assayed through the colorimetric method at 570 nm using a Thermo Scientific Evolution 600 UV-Vis spectrophotometer and the results are expressed as $\text{mmoles H}_2\text{O}_2/\text{min}/\text{g}$ at 25°C (Babeanu et al., 2010).

Proline content was determined in 3% aqueous sulfosalicylic acid extract by spectrophotometry at 520 nm following the ninhydrin method, using L-proline as a standard. The results are expressed as μg proline/ g fw (Matei et al., 2013).

All assays were performed in triplicate and the results presented here are the mean values.

RESULTS AND DISCUSSION

The plants grown on ash deposits are subject to several stress factors: heavy metals, lack of nutrients and water, heat, etc. The responses to these types of stress depend on the species and genotype, the developmental and metabolic state of the plant and the duration and severity of the stress (Singh and Tuteja, 2010).

The results for peroxidase activity are shown in Figure 1. Peroxidase activity varies between 4.00 $\Delta A/\text{min}/1\text{g}$ f.w. (Pesred) and 10.26 $\Delta A/\text{min}/1\text{g}$ f.w. (Fragisal). Swedish *Salix* clones showing the lowest average value for peroxidase activities: (6.025 $\Delta A/\text{min}/1\text{g}$ fw) while Romanian clones show higher values (7.53 $\Delta A/\text{min}/1\text{g}$ fw).

In a previous research paper the same willow clones were investigated in a control plantation at Radovan area with well-developed plants and a plantation on sandy soil at Tamburesti. For the control plantation Radovan, with plants well hydrated values obtained for peroxidase activity varies between 1.62 $\Delta A/\text{min}/1\text{g}$ f.w.-Inger and 15.38 $\Delta A/\text{min}/1\text{g}$ f.w.-RO1082 (Babeanu et al., 2017).

In the case of plants grown on fly ash the peroxidase activity increases compared to control plantation Radovan (Pesred: 22.32% -Olof: 3.23 fold) with two exception: RO 892 and RO 1082 which had a decrease in enzymatic activity of peroxidase.

Our results are in agreement with other studies reporting the increased peroxidase activity in response to water stress and heavy metal stress in other plants (Babeanu et al., 2010). Increasing peroxidase activity leads to lower H₂O₂ content in the cell and to avoid oxidative stress.





Figure 1 Peroxidase activity in leaves of Salix clones

The obtained results for catalase activity are shown in Figure 2. In the case of plants grown on fly ash deposit catalase activity varies between 683.5 μM H₂O₂/min/g (Olof) and 885.3 μM H₂O₂/min/g (RO1082). In this case catalase activity increases compared to control plantation Radovan (from 2.72 % -Olof to 73.84 % -RO 1077) except for the clones Pesred, Cozia, Tordis, Jorr and Torhild which had a decrease in catalase activity (8.83%, 12.09%, 11.22 %, 11.56% respectively 3.28%).

For the control plantation Radovan, catalase activity ranges from 423.53 μM H₂O₂/min/g - RO1077 to 947.26 μM H₂O₂/min/g –Jorr (Babeanu et al., 2017).


Figure 2 Catalase activity in leaves of Salix clones



The proline content varies with analyzed clone (figure 3). In the case of plants grown on fly ash the proline content varies between 198.16 μg/1g fw (Cozia) and 414.75 μg/1g fw (Tora).

The proline content for all investigated clones is higher in plants grown on fly ash compared to reported data for well-developed plants grown in the Radovan plantation.

Figure 3 Proline content in leaves of studied Salix clones



For the control plantation Radovan, with plants well hydrated, the results obtained for proline content varies between 95.61 µg/1g fw-RO1077 and 210 µg/1g fw-RO1082 (Babeanu et al., 2017). Accumulation of proline as a signal for adaptive plant responses to stress has been reported for many species (Babeanu et al., 2012, Storaska et al., 2008).

CONCLUSIONS

The analyzed biochemical indices show a dependency with the investigated genotypes

Fly ash conditions may disturb the redox homeostasis and lead to oxidative stress, increasing production of reactive oxygen species. Balance between the production and the scavenging of reactive oxygen species is critical to the maintenance of growth and metabolism of plants.

The high levels of peroxidase and catalase activity and proline content suggests a state of oxidative stress, the plants activating a defensive system.

ACKNOWLEDGEMENTS

This paper was financially supported by MEN UEFISCDI, Programme PN II 2014-2016, project no. 111 SAROSWE.

BIBLIOGRAPHY

1. **Babeanu C., Constantinescu E., Popa D., 2004.** *Effect of nitrogen and phosphorus fertilizers on the agrochemical properties of ash crops-from Isalnita power plant-cultivated with corn, Journal of Environmental Protection and Ecology, 5 (2) 403-408*
2. **Babeanu C., Constantin C., Paunescu G., Popa D., 2010.** *Effects of drought stress on some oxidoreductase enzymes in five varieties of wheat, Journal of Environmental Protection and Ecology 11 (4) 1280-1284*
3. **Babeanu C., Păunescu A., Matei Ghe., 2012.** *Free amino acids and proline accumulation in wheat leaves under water stress, Annals of the University of Craiova, The Chemistry Series, XLI (2) 23-26*

4. **Babeanu C., Soare M., Corneanu M., Dragoi M.,** 2017. *Effects of drought stress on some oxidoreductase enzymes and proline content in leaves of Salix genotypes*, SGEM2017 Conference Proceedings, Vol. 17, Issue 61, 665-672
5. **Hernea C., Trava I.D., Borlea Gh. F.,** 2015. *Biomass production of some Swedish willow hybrids on the West of Romania. A case study*, Journal of Horticulture, Forestry and Biotechnology, 19 (3) 103- 106
6. **Hernea C., Corneanu M., Sărac I., Turcu E.,** 2016. *The potential of romanian and swedish willow for short rotation coppice in the specific condition of Banat plain: a comparative analyze*, South west J Hortic Biol Environ, 7(2), 115-126
7. **Ibrahim I.A.A,** 2015. *Chemical characterization and mobility of metal species in fly ash–water system*, Water Science 29, 109-122
8. **Matei Ghe., Babeanu C., Soare M., Birzanu O.F.,** 2013. *The effect of cadmium excess on lipid peroxidation and antioxidant defence system in wheat seedlings*, Analele Universității din Craiova, seria Agricultură – Montanologie – Cadastru (Annals of the University of Craiova - Agriculture, Montanology, Cadastre Series) 43 (1) 211-216
9. **Popa D., Hanescu V., Constantin C., Codreanu M.,** 2010. *The concept of plant-fungi-soil partnership- a realistic approach of the symbiotic systems developed on the dumps of ashes from Isalnita area-Romania*, Journal of Environmental Protection and Ecology, 11 (3), 958-965
10. **Popa D., Codreanu M.D., Babeanu C.,** 2013. *Mycorrhizal colonisation – a possible alternative biotechnology recommended to the fly ashes dumps*, Journal of Environmental Protection and Ecology, 14 (1), 256-262
11. **Singh G. S., Tuteja N.,** 2010. *Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants*, Plant Physiology and Biochemistry, 48, 909-930
12. **Soare M., Păniță O., Sălceanu C.,** 2015. *Partial results concerning the behavior of energy willow genotypes in cultivated improper areas*, Annals of the University of Craiova: Agriculture, Montanology, Cadastre Series, XLV, 300-305
13. **Soare M., Iancu P., Panita O., Soare R., Bonciu E.,** 2017. *Researches concerning the possibility of cultivating energetic willow on deposit of ash from thermal power stations*, SGEM2017 Conference Proceedings, Vol. 17, Issue 42, 519-528
14. **Soare R., Soare M., Rosculete E., Panita O.,** 2010. *Preliminary research concerning the polluted substances contain from the soil and water inside of some conventional vegetable farms from South Plain of Romania*, Annals of the University of Craiova- Agriculture, Montanology, Cadastre Series. 40(2) 277-282
15. **Stolarska A., Wróbel J., Przybulewska K.,** 2008. *Free proline content in leaves of Salix viminalis as an indicator of their resistance to substrate salinity*, Ecological Chemistry And Engineering, 15 (1-2), 139-146